

blank page

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## Triclopyr/Picloram 400 EC SELECTIVE HERBICIDE

### ACTIVE CONSTITUENTS:

300 g/L TRICLOPYR present as BUTOXYETHYL ESTER

100 g/L PICLORAM present as HEXYLOXYPROPYLAMINE SALT

GROUP **4** HERBICIDE

For control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: Read this leaflet before using this product**

### AgMerch Pty Ltd

ABN 26 645 371 017

217 Wyndham Street,  
Shepparton VIC 3630

m: 0498 530 214

info@agmerch.com.au

www.agmerch.com.au

APVMA Number: 92094/134487

## DIRECTIONS FOR USE

### RESTRAINTS:

**DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

**DO NOT** spray if rain is likely within one hour or if foliage is wet from rain or dew.

**DO NOT** burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

**DO NOT** apply by aerial application in wind in excess of 15 km/hr and air temperatures above 35°C.

In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

### 1. WOODY WEED SITUATIONS

#### Table A: High Volume Spraying

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>			
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/100 L WATER</b>
African boxthorn	Less than 2 m tall	All States	500 mL
<i>Angophora</i> spp.	1 to 3 m tall		350 mL
Australian blackthorn	Less than 2 m tall		500 mL
<i>Banksia</i> spp.	1 to 3 m tall		350 mL
Biddy Bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	ACT and NSW only	500 mL
Blackberry in association with: Docks, Ragwort, Smartweed, Thistles	Late spring to autumn	All States except NT	350 mL OR 500 mL
Blue heliotrope	Flowering	NSW and Qld only	500 mL
Brooms: Cape, English, Flax leaf, Montpellier	Spring to mid-summer prior to pod formation	All States except NT	250 mL
	Autumn to winter		350 mL
Camphor laurel	Less than 2m tall	All States	
	Above 2 m tall		500 mL
<i>Casuarina</i> spp.	1 to 3 m tall		350 mL
Chinese apple	Less than 2 m tall	Qld and WA only	
Cockspur thorn	Spring to autumn	ACT, NSW and Qld only	
Common sensitive plant	Any time when actively growing	NT, Qld and WA only	200 mL

<b>CRITICAL COMMENTS</b>
Apply when bushes have good leaf cover, growth and no leaf fall.
Apply from late spring to early autumn.
Add a 100% concentrate non-ionic surfactant at 125 mL/100 L of water for best results.
<b>Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry.</b> Where herbicides other than Group 4 Herbicides have been used, allow two seasons regrowth to occur before respraying with AgMerch Triclopyr/Picloram 400 EC Selective Herbicide. Apply in a minimum spray volume of 1250 L/ha.
Apply as a thorough foliage spray.
Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Apply as a thorough foliage spray.
To avoid leaves closing during application, spray plants while moving forward. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.**

<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/100 L WATER</b>
Crofton weed	Spring to autumn	ACT, NSW and Qld only	350 mL
Eastern cotton bush ( <i>Maireana microphylla</i> )	Spring to autumn	NSW and Qld only	500 mL
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3 m tall	All States	350 mL OR 500 mL
Galenia	Fresh growth in spring to summer	NSW only	500 mL
Giant bramble	Spring to autumn	NT, Qld and WA only	
Gorse	1 to 1.5 m tall	All States except NT	250 mL
	Over 1.5 m tall or autumn treatment		350 mL
	Winter treatment		500 mL
Groundsel bush ( <i>Baccharis halimifolia</i> )	1 to 1.5 m tall in spring and summer	All States except NT	250 mL
	Over 1.5 m tall or autumn treatment		350 mL
Green cestrum	Late spring to early autumn	ACT, NSW and Qld only	500 mL
Hawthorn	Less than 2 m tall	All States	
Horehound	Pre-flowering	NSW and Qld only	350 mL
Japanese sunflower			
Lantana ( <i>Lantana camera</i> ) ( <i>Lantana montevidensis</i> )	Up to 1 m tall in summer to autumn	All States	
	1 to 2 m tall in summer to autumn		500 or 750 mL
Lion tail ( <i>Leonatis nepetifolia</i> )	Pre-flowering	Qld only	200 mL
Limebush	Any time of year with good leaf cover and soil moisture	NT and Qld only	350 mL

<b>CRITICAL COMMENTS</b>
Apply as a thorough foliage spray.
Add Uptake® Spraying Oil at 500 mL/100 L of water. Some bushes may require a follow-up spray to control regrowth. <b>Apply the high rate where difficult to control species of Eucalyptus regrowth is present.</b> Addition of an adjuvant may improve results – contact AgMerch Pty Ltd for details.
Use 2000 L of spray mixture/ha.
Penetration of thick clumps may be difficult, and respraying may be necessary. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results. Spring and Summer treatment only. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Brownout may not be complete until summer. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results. Apply as thorough foliar spray.
One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high. Apply from late spring to early autumn. Apply as a thorough foliar spray.
Add one of the following adjuvants, when using 350 mL rate: Uptake® Spraying Oil @ 0.5% v/v. Pulse® @ 0.1% v/v. Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Apply as a thorough foliar spray. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results – contact AgMerch Pty Ltd for details.

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.**

<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/100 L WATER</b>
Manuka	At flowering	Vic only	500 mL
Mesquite ( <i>Prosopis</i> spp.)	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350 mL
<i>Prosopis velutina</i>		Qld only	670 mL
Mistflower	Spring to autumn	ACT, NSW and Qld only	
Mother-of-millions	Flowering	NSW and Qld only	500 mL
Paddy's lucerne	Active growth	NSW only	
Parkinsonia	Under 2 m tall	NT, Qld and WA only	350 mL
Prickly Pear (common) Smooth tree pear	Active phyllode growth	All States	500 mL
Rubber vine (Not infected with rust)	Up to 1.5 m tall at flowering	NT, Qld and WA only	350 mL
	Dense stands greater than 1.5 m tall at flowering		500 mL
Siam weed	Active growth	Qld and WA only	350 mL
Sicklepod	Up to flowering	NT, Qld and WA only	200 mL
St John's wort	From flowering to early seed set	All States	500 mL
Sweet briar	Up to 1.5 m tall	All States except NT	350 mL
			500 mL
Tobacco weed	Actively growing plants	NT, Qld and WA only	300 mL
Wattle ( <i>Acacia</i> spp.) (except corkwood wattle)	1 to 3 m tall	All States	350 mL
Wild rosemary ( <i>Cassinia laevis</i> )	Active growth, 0.5 to 1 m tall	Qld only	350 to 500 mL
Wild tobacco tree	Spring to autumn up to 2 m tall	ACT, NSW and Qld only	350 mL

<b>CRITICAL COMMENTS</b>
For optimum results, add a registered adjuvant containing 1020 g/L Polyether modified polysiloxane at 200 mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants. <b>DO NOT</b> spray plants bearing pods. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Apply as a thorough foliar spray.
Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur. Add Uptake® Spraying Oil at 500 mL/100 L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage. Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands, regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Late spring to early summer. Add Metsulfuron 600 WG Herbicide at 10 g/100 L water to obtain more reliable results with the lower rate of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water for best results.
Use lower rate on seedlings 0.5 m tall. Apply as a thorough foliar spray.

**Table B: Aerial Application**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.**

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100 L WATER
Blackberry	Summer to autumn	NSW, Qld, SA, Vic and WA only	10 L
Goose		Tas only	
Cockspur thorn, Crofton weed, Lantana, Mistflower	Late autumn	NSW, NT and Qld only (helicopter only)	1.5 L plus 7.5 L 2,4-D amine (500 g/L formulation)
Lantana			10 L
Rubber vine (Not infected with rust)	When flowering	NT and Qld only (helicopter only)	3 L to 5 L
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4 L

**AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS**

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100 L WATER
Parkinsonia	Seedlings, 1 - 2 m tall or 12 to 24 months old	Qld and NT only (helicopter only)	3 L

**Table C: Controlled Droplet Application (C.D.A.)**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.**

WEEDS CONTROLLED	WEED GROWTH CONTROLLED	STATE	RATE/ha
Blackberry in association with: Docks, Ragwort, St John's wort, Thistles	Summer to autumn	All States except NT	Apply undiluted.

CRITICAL COMMENTS
Where herbicides other than Group 4 Herbicides have been used, allow two seasons regrowth to occur before respraying with AgMerch Triclopyr/Picloram 400 EC Selective Herbicide.
<b>WARNING:</b> Eucalyptus species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover.
Helicopter application only.
Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha.
<i>Follow-up respraying will be required.</i>
Helicopter application only.
Use rates will depend upon the density and height of the rubber vine stand. The higher rates should be used on dense stands; however, complete coverage and penetration may be difficult.
<i>Follow-up respraying will be required.</i>
Any regrowth should be sprayed with a suitable basal bark herbicide.
Helicopter application only.
<i>Follow-up spraying will be required in the following season.</i>

CRITICAL COMMENTS
Add Uptake® Spraying Oil at 1 L/ha.

CRITICAL COMMENTS
One application may give satisfactory control, but subsequent regrowth and seedlings should be re-sprayed after hardening off. Where herbicides other than Group 4 Herbicides have been used, allow two seasons regrowth to occur before respraying with AgMerch Triclopyr/Picloram 400 EC Selective Herbicide.

**Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer)**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.**

WEEDS CONTROLLED	WEED GROWTH CONTROLLED	STATE	RATE/ha
Blackberry	Late spring to autumn	ACT, NSW, Qld, SA, Tas and WA only	335 mL
Camphor laurel, Cockspur thorn, Crofton weed	Less than 1.5 m high	ACT, NSW and Qld only	500 mL
Eucalyptus species	Seedlings up to 2 m tall	All States	335 mL
Mistflower	Less than 1.5 m high	ACT, NSW and Qld only	500 mL
Sweet briar	1.5 m tall, full leaf to ripe fruit	NSW only	
St John's wort	During flowering to early seed set	NSW, Tas and Vic only	
Wild tobacco tree	Less than 1.5 m high	ACT, NSW and Qld only	

**Table E: Boom Application**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

**AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.**

WEEDS CONTROLLED	WEED GROWTH CONTROLLED	STATE	RATE/ha
Galenia	Fresh growth during spring to summer	NSW only	5 L
Sicklepod	Up to flowering	NT and Qld only	3 L
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4 L

CRITICAL COMMENTS
Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high-volume application technique is recommended.
<b>Gas Powered Gun only:</b> Apply to actively growing bushes not more than 1.5 m tall that have not more than 5 stems from the crown.
<b>Gas Powered Gun only:</b> One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high-volume application technique is recommended.

CRITICAL COMMENTS
Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100 mL/100 L of water. Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

## 2. FALLOW SITUATIONS

**Table A: Boom Application**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha
Blackberry nightshade – Suppression only	10 to 25 cm tall, prior to flowering	NSW and Qld only	200 to 400 mL + 1.2 L Glyphosate 450 Herbicide + Adjuvant
Camel melon, Prickly paddy melon, Cucumber melon ( <i>Cucumis melo</i> )	From 2 leaf to 50 cm diameter		
Common sowthistle	From 8 leaf to flowering		
Cow vine	From 2 to 5 leaf up to 15 cm diameter, prior to flowering		
Lucerne (established)	Active growth, 15 to 25 cm high, during spring		
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20 cm diameter, prior to flowering	300 to 500 mL + 1.2 L Glyphosate 450 Herbicide + Adjuvant	200 to 400 mL + 1.2 L Glyphosate 450 Herbicide + Adjuvant

**Table B: Blanket Wiper Application**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW			
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha
Bitterbark ( <i>Alstonia constricta</i> )	From summer to end of autumn	Qld	1:4 (1 part AgMerch Triclopyr/Picloram 400 EC Selective Herbicide to 4 parts water) 2% solution for spot spray (e.g. 100 mL AgMerch Triclopyr/Picloram 400 EC Selective Herbicide in 5 L water)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY: DO NOT treat bushes carrying mature or near mature fruit.**

**FOR NATIVE VEGETATION:** Use of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**

### CRITICAL COMMENTS

#### FOR USE BY GROUND EQUIPMENT ONLY.

Plants must be actively growing.

Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for *Polymeria pusilla*).

Refer to Glyphosate 450 Herbicide label for use of adjuvant.

**DO NOT** plant susceptible crops for up to 9 months after application, as specified in General Instructions – Minimum Recropping Periods – Black Cracking Clay Soils, NSW & Qld.

Dry conditions after application will increase the recropping interval.

### CRITICAL COMMENTS

For use with blanket wipers only.

For best results apply in autumn to tall (> 60 cm) plants using two opposite directional passes (up and back). Follow up “missed” plants with a spot spray application. These will be obvious after 6 weeks.

Blanket wiper applications can be made in summer when plants are smaller but follow up spot spraying may be necessary.

Do not disturb (cultivate) the treated patches for at least 3 months. Best long-term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).

Spot spraying “missed” plants:

Thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.

## GENERAL INSTRUCTIONS

### MINIMUM RE-CROPPING PERIODS – Black Cracking Clay Soils, NNSW & Qld.

**Table A: Boom Application**

Plant-back periods for crops following the application of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide for rates up to 600 mL/ha				
RATE mL/ha	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plant back period may be significantly longer.

**Table B: Blanket Wiper Application**

Plant-back periods for crops following blanket wiper application	
CROP	Months
Broadleaf crops	18
Lucerne	6
This will allow any potential soil residues to dissipate, if any, and allow effective control.	

**Note:** Before using AgMerch Triclopyr/Picloram 400 EC Selective Herbicide in tank mixes with other herbicides, check the plant back information on all product labels. The most residual product, i.e. the product with the longest plant-back period, will determine the time between spraying and planting.

## GENERAL INSTRUCTIONS

### COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with AgMerch Triclopyr/Picloram 400 EC Selective Herbicide (e.g. 2,4-D amine).

AgMerch Triclopyr/Picloram 400 EC Selective Herbicide is compatible with the following herbicides:

2,4-D, Metsulfuron, Glyphosate 450 Herbicide, Fluroxypyr, Glyphosate 490.

AgMerch Triclopyr/Picloram 400 EC Selective Herbicide is compatible with the following adjuvants, as per Directions for Use:

Uptake® Spraying Oil, a registered adjuvant containing 1020 g/L Polyether modified polysiloxane and non-ionic surfactant (1000 g/L).

## MIXING

Mix only with water.

Half fill the spray unit with water, and add the required amount of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

## APPLICATION

### 1. WOODY WEED SITUATIONS

**Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.**

#### A. High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

##### Hand Gun

- Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's wort).
- A spray volume of 3000 to 4000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100 m<sup>2</sup>) should be used.
- Use 2000 L of spray mixture/ha of Galenia infestation (i.e. 20 L/100 m<sup>2</sup> infested area).

##### Knapsack

- Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun.
- A spray volume of 3 to 4 L/10 m<sup>2</sup> infested area should be used.
- A spray volume of 2 L/10 m<sup>2</sup> should be used for an area infested with Galenia.

#### B. Aerial Application

- Apply in 200 L of water/ha using an aircraft to apply 100 L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter.
- The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.
- DO NOT spray when wind exceeds 15 km/hr and/or air temperature reaches 35°C.

#### C. Controlled Droplet Application (C.D.A.)

- Results similar to high volume spraying can be obtained using Micron Herbi\* or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided by C.D.A. unit.

#### D. Low Volume High Concentrate Application Techniques

- Good control will be achieved, similar to high volume application, where bush size enables good



coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

- **Gas Powered Gun:** Apply 50 mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.
- **Sprinkler sprayer:** This technique involves using a micro sprinkler that is connected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

### E. Boom Application

- Application in a minimum spray volume of 200 L/ha for Galenia and St John's wort and 600 L of water/ha for Sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

## 2. FALLOW SITUATIONS

### A. Boom Application

Application of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

### B. Blanket Wiper Application

- Blanket needs to be made from durable and wettable material with a rigid backing.
- Blanket should be rigidly mounted behind motorised vehicle (e.g. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting).
- Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle.
- Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15 kph are ideal for blanket wiping application.

## CLEANING SPRAY EQUIPMENT

### Rinsing

- After using AgMerch Triclopyr/Picloram 400 EC Selective Herbicide, empty spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.
- After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Decontamination

- Before spraying cotton and other sensitive crops with equipment that has been used to apply AgMerch Triclopyr/Picloram 400 EC Selective Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

- Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF®, OMO®, OMOMATIC®, DRIVE® at 500 mL/100 L of water or the powder equivalent at 500 g/ 100 L of water) and circulate throughout the system for at least 15 minutes.
- Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

## RESISTANT WEEDS WARNING



AgMerch Triclopyr/Picloram 400 EC Selective Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group 4 Herbicide.

Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist though normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 4 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AgMerch Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local AgMerch Pty Ltd representative.

## PRECAUTIONS

### Re-Entry

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to AgMerch Triclopyr/Picloram 400 EC Selective Herbicide include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and Pinus spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines.

AgMerch Triclopyr/Picloram 400 EC Selective Herbicide is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product can remain in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.

DO NOT apply close to, or in areas containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants.

DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.

### **PROTECTION OF LIVESTOCK**

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilizers or seed.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length chemical resistant gloves and face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia:13 11 26.

### **SAFETY DATA SHEET**

For further information refer to Safety Data Sheet (SDS) which can be obtained from the supplier.

### **CONDITIONS OF SALE**

The use of AgMerch Triclopyr/Picloram 400 EC Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by AgMerch Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AgMerch Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.